

Underlined parts: revised in July 2019

[Keep this information leaflet with this medication and be sure to read it before taking the medication.]

NEW SSBRON® TABLETS ACE

Class ② OTC drug

Antitussive-expectorant medication containing L-carbocysteine, a mucolytic agent, for persistent cough and sputum

- **NEW BRON TABLETS ACE** exert excellent effects against cough with sputum, by the actions of the following ingredients of the tablets: ingredients to alleviate cough, promote the expectoration of sputum stuck in the throat, and alleviate respiratory distress.

Precautions for Use



Contraindications

(If you do not follow the precautions exactly as listed below, your current symptoms may worsen, and you may be placed at increased risk of side effects or accidents.)

- ❶ **The following persons must not take this medication:**
 - (1) Persons who have had an allergic reaction to any of the ingredients of NEW BRON TABLETS ACE
 - (2) Children under 12 years of age
- ❷ **Do not use any of the following medications during the period of taking NEW BRON TABLETS ACE:**

Other antitussive-expectorant medications, common cold medications, sedatives, and oral medications containing antihistamines (e.g., oral rhinitis medications, motion sickness medications, anti-allergic medications, and hypnotic sedatives)
- ❸ **Do not operate vehicles or machinery after taking this medication.**

(Sleepiness or drowsiness may occur.)
- ❹ **Breast-feeding women are advised to not take NEW BRON TABLETS ACE or stop breast-feeding if this medication is to be taken.**
- ❺ **Do not take more than the specified dosage of this medication or use it over a long continuing period.**



Consultation

- ❶ **The following persons should consult a physician, pharmacist, or registered salesperson before taking this medication:**
 - (1) Persons who are receiving any treatment from a physician
 - (2) Women who are or may be pregnant
 - (3) Older persons
 - (4) Persons who have had any allergic reactions to drugs or other substances
 - (5) Persons with the following symptom: Hyperthermia (high fever) and dysuria
 - (6) Persons who have received a diagnosis of any of the following diseases: heart diseases, hypertension, diabetes mellitus, glaucoma, liver diseases, thyroid dysfunction, respiratory dysfunction, obstructive sleep apnea syndrome, or obesity
- ❷ **If any of the symptoms listed below develop after taking NEW BRON TABLETS ACE, they might be side effects of the medication. Stop taking this medication immediately and consult a physician, pharmacist, or registered salesperson, presenting this leaflet.**

Affected area	Symptoms
Skin	Rash/redness and itching
Gastrointestinal	Abdominal pain, nausea/vomiting, or inappetence
Neuropsychiatric	Dizziness
Urinary	Dysuria

In rare cases, any of the serious symptoms listed below may occur. In such cases, immediately consult a physician.

Name of the symptoms	Symptoms
Shock (anaphylaxis)	Itchy skin, urticaria, hoarse voice, sneezing, itchy throat, respiratory distress, palpitations, clouding of consciousness, etc., occurring immediately after taking this medication.
Oculomucocutaneous syndrome (Stevens-Johnson syndrome) or toxic epidermal necrolysis	Persistence or sudden worsening of symptoms such as high fever, ocular hyperemia, eye discharge, lip erosion, sore throat, or rashes/redness over an extensive area of the skin
Hepatic impairment	Occurrence of fever, itching, rash, jaundice (yellowing of the skin and whites of the eyes), brown urine, general malaise, inappetence, etc.
Aplastic anemia	The following symptoms can occur: bruising, nasal bleeding, gingival bleeding, fever, pallor of the skin or mucous membranes, fatigue, palpitations, shortness of breath, feeling sick and fainting, and hematuria.
Agranulocytosis	Occurrence of sudden high fever, chills, sore throat, etc.
Respiratory depression	Occurrence of shortness of breath, respiratory distress, etc.

(Continued from the front side)

③ **Any of the conditions mentioned below may occur after taking NEW BRON TABLETS ACE. If such symptoms persist or worsen, stop taking this medication and consult a physician, pharmacist, or registered salesperson, presenting this leaflet.**

Constipation, thirst, sleepiness, or diarrhea

④ **If the symptoms for which this medication is being taken are not alleviated after 5 or 6 doses, stop taking this medication and consult a physician, pharmacist, or registered salesperson, presenting this leaflet.**

Indications Cough and sputum

Dosage and Administration Take a dose of NEW BRON TABLETS ACE with cold or lukewarm water thrice daily after meals, according to the table below.

Age	Adults (aged 15 years or older)	Children aged 12 to 14 years	Children under 12 years of age
Dose	4 tablets	2 tablets	×This medication must not be taken.

< Precautions Concerning Dosage and Administration >

- (1) NEW BRON TABLETS ACE must be used strictly in accordance with the specified dosage and administration.
- (2) Children must take this product only under the supervision of their parents/guardians.

Ingredients and Actions

Ingredients	Amount per 12 tablets	Action
L-Carbocisteine	750 mg	To reduce the viscosity (thickness) of sputum stuck in the throat and normalize the respiratory mucosa damaged by coughing, thereby promoting sputum expectoration
Dihydrocodeine phosphate	30 mg	To inhibit the cough reflex by acting on the cough center
dl-Methylephedrine hydrochloride	75 mg	To relieve bronchial muscle tone and eliminate respiratory distress, alleviating cough
Chlorpheniramine maleate	12 mg	To alleviate allergy cough

Excipients: carmellose calcium, silicic anhydride, cellulose, lactose, white soft sugar, hydroxypropylcellulose, hypromellose, povidone, calcium carbonate, carnauba wax, magnesium stearate, talc, titanium oxide, pullulan, and polyoxyethylene polyoxypropylene glycol


For the treatment of cough and sputum...

In addition to taking this medication, attention should be paid to the following points:

- Try to restore physical strength by following a nutritious diet and having sufficient sleep.
- Make sure indoor air does not become overly dry.
- Do not smoke and pay attention to throat care such as by gargling.

Precautions for Storage and Handling

- (1) Store the product in the jar with the lid closed tightly, in a dry, cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer the contents to another container (which may lead to drug misuse or changes in the quality of this product).
- (4) Do not handle this product with wet hands. If the product is handled with wet hands, the sugar coating may become uneven or discolored.
- (5) The filler in the jar is intended to prevent breakage of tablets during transportation. After the package has been opened, the filler is no longer necessary and must be removed.
- (6) Do not take this medication after the expiration date.



Contact Information

Inquiries should be made to the drugstore from where you purchased the product or to the customer service office.
Customer Service Office, SSP Co., Ltd. Phone: 0120-028-193
Opening hours: 9:00 to 17:00 (closed on Saturdays, Sundays, and Japanese public holidays)



Manufactured and Distributed by:

SSP Co., Ltd.

3-20-2 Nishi-Shinjuku, Shinjuku-ku, Tokyo 163-1444
<https://www.ssp.co.jp/>