

S.TAC® GENERAL COLD MEDICINE

Class ② OTC drug

General cold medicine: Film-coated tablets for common cold that can be taken by family members of all ages (5 years and above)

- S.TAC GENERAL COLD MEDICINE is a medication for common cold that can be taken by family members of all ages (5 years and above).
- This medication is an easy-to-swallow, film-coated, tablet formulation that is designed to work effectively on 11 types of common cold symptoms.



Precautions for Use



Contraindications

If you do not follow the precautions exactly as listed below, your current symptoms may worsen, and you may be placed at increased risk of side effects or accidents.

① **The following persons must not take this medication:**

- (1) Persons who have had an allergic reaction to any of the ingredients of S.TAC GENERAL COLD MEDICINE
- (2) Persons who have experienced asthma after taking this medication, other common cold medications, or antipyretic-analgesics

② **Do not use any of the following medications during the period of taking S.TAC GENERAL COLD MEDICINE:**

Other common cold medications, antipyretic-analgesic medications, sedatives, antitussive-expectorant medications, or oral medications containing antihistamines (e.g., oral rhinitis medications, motion sickness medications, anti-allergic medications, or hypnotic sedatives)

③ **Do not operate vehicles or machinery after taking this medication.**

(Sleepiness or drowsiness may occur.)

④ **This medication must not be taken immediately before or after alcohol intake.**

⑤ **Do not use this medication over a long continuing period.**



Consultation



① **The following persons should consult a physician, pharmacist, or registered salesperson before taking this medication:**

- (1) Persons who are receiving any treatment from a physician or dentist
- (2) Women who are or may be pregnant
- (3) Breast-feeding women
- (4) Older persons
- (5) Persons who have had any allergic reactions to drugs or other substances
- (6) Persons with the following symptom: Hyperthermia (high fever) and dysuria
- (7) Persons who have received a diagnosis of any of the following diseases: thyroid dysfunction, diabetes mellitus, heart disease, hypertension, liver disease, kidney disease, gastroduodenal ulcers, or glaucoma

② **If any of the symptoms listed below develop after taking S.TAC GENERAL COLD MEDICINE, they might be side effects of the medication. Stop taking this medication immediately and consult a physician, pharmacist, or registered salesperson, presenting this leaflet.**

Affected area	Symptoms
Skin	Rash/redness and itching
Gastrointestinal	Nausea/vomiting and inappetence
Neuropsychiatric	Dizziness

Affected area	Symptoms
Respiratory	Shortness of breath or respiratory distress
Urinary	Dysuria
Other	Excessively decreased body temperature

In rare cases, any of the serious symptoms listed below may occur. In such cases, immediately consult a physician.

Name of the symptoms	Symptoms
Shock (anaphylaxis)	Itchy skin, urticaria, hoarse voice, sneezing, itchy throat, respiratory distress, palpitations, clouding of consciousness, etc., occurring immediately after taking this medication
Oculomucocutaneous syndrome (Stevens-Johnson syndrome), toxic epidermal necrolysis, or acute exanthematous pustulosis	Persistence or worsening of the following symptoms: high fever, ocular hyperemia, eye discharge, lip erosion, sore throat, rashes/redness over an extensive area of the skin, small rashes (small pustules) appearing on reddened skin, general malaise, or inappetence
Drug-induced hypersensitivity syndrome	Occurrence of symptoms such as redness over an extensive area of the skin, generalized exanthema, fever, malaise, or swelling of the lymph nodes (cervical, axillary, and/or inguinal lymph nodes)
Hepatic impairment	Occurrence of fever, itching, rash, jaundice (yellowing of the skin and whites of the eyes), brown urine, general malaise, inappetence, etc.

(Continued on the reverse side)

®: Registered trademark

(Continued from the front side)

Name of the symptoms	Symptoms
Renal disorders	Occurrence of fever, rashes, decreased urine output, generalized edema/swelling, general malaise, arthralgia (pain in joints), diarrhea, etc.
Interstitial pneumonia	Sudden occurrence or persistence of the following symptoms: shortness of breath or respiratory distress when climbing stairs or slightly overworking, dry cough, or fever
Asthma	Occurrence of breathing sounds, such as wheezing and whistling, or respiratory distress
Aplastic anemia	The following symptoms can occur: bruising, nasal bleeding, gingival bleeding, fever, pallor of the skin or mucous membranes, fatigue, palpitations, shortness of breath, feeling sick and fainting, and hematuria.
Agranulocytosis	Sudden high fever, chills, sore throat, etc.

③ Any of the conditions mentioned below may occur after taking S.TAC GENERAL COLD MEDICINE. If such symptoms persist or worsen, stop taking this medication and consult a physician, pharmacist, or registered salesperson, presenting this leaflet.
Thirst and sleepiness

④ If the symptoms of common cold are not alleviated after taking 5 or 6 doses, stop taking this medication and consult a physician, pharmacist, or registered salesperson, presenting this leaflet.

Indications

Alleviation of various symptoms of common cold (sore throat, nasal discharge, nasal stuffiness, fever, chills, headache, sneezing, cough, sputum, joint pain, and muscle pain)

Dosage and Administration

Take a dose of S.TAC GENERAL COLD MEDICINE, according to the table below, with cold or lukewarm water, 3 times daily. Each dose should be taken within 30 minutes after a meal, whenever possible.

Age	Adults (aged 15 years or older)	Children aged 11 to 14 years	Children aged 5 to 10 years	Children aged under 5 years
Dose	 3 tablets	 2 tablets	 1 tablet	× This medication must not be taken.

< Precautions Concerning Dosage and Administration >

(1) S.TAC GENERAL COLD MEDICINE must be used strictly in accordance with the specified dosage and administration.

(2) Children must take this product only under the supervision of their parents/guardians.

Acetaminophen	900 mg	Kanzo (glycyrrhiza) extract
Chlorpheniramine maleate	7.5 mg	(equivalent to 750 mg of glycyrrhiza)
Dextromethorphan		187.5 mg
hydrobromide hydrate	48 mg	Powdered ginger
DL-methylephedrine hydrochloride	60 mg	Anhydrous caffeine
Hesperidin	45 mg	75 mg

Excipients: croscarmellose sodium, silicic anhydride, cellulose, lactose, hypromellose, povidone, macrogol, magnesium stearate, talc, titanium oxide, corn starch, and red ferric oxide

Precautions for Storage and Handling

- Store in a dry, cool place away from direct sunlight.
- Keep out of the reach of children.
- Do not transfer the contents to another container (which may lead to drug misuse or changes in the quality of this product).
- Close the lid of the jar tightly after each use. If the lid of the jar is not tightly closed, the product may deteriorate due to the effects of moisture, etc. Do not handle this product with wet hands.
- The filler in the jar is intended to prevent breakage of tablets during transportation. After the package has been opened, the filler is no longer necessary and must be removed.
- Do not take this medication after the expiration date.



Contact Information

Inquiries should be made to the drugstore from where you purchased the product or to the customer service office.
Customer Service Office, SSP Co., Ltd. Phone: 0120-028-193
Opening hours: 9:00 to 17:00 (closed on Saturdays, Sundays, and Japanese public holidays)



Manufactured and Distributed by:
SSP Co., Ltd.

3-20-2 Nishi-Shinjuku, Shinjuku-ku, Tokyo 163-1444
<https://www.ssp.co.jp/>

