

この説明書は製品についての参考情報です。正しくは製品のパッケージをご確認ください。パッケージデザインは変更になっている可能性があります。

This document serves as reference information. Please ensure to read the information on the package.

The package design may have been updated

[Precautions for Use] Contraindications (If you do not follow the precautions exactly as listed below, your current symptoms may worsen and you may be placed at increased risk of side effects.) **1. The following persons must not take this product:** persons who (1) have hyperchlorhydria (excessive gastric acid secretion) and (2) have received a diagnosis of heart disease or gastric ulcers. **2. Do not take the following medications during the period of taking ESTARON MOCHA ORAL SOLUTION:** other prophylactic medications against sleepiness. **3. Do not take this product with caffeine-containing beverages such as coffee and tea.** **4. Take ESTARON MOCHA only over a short period of time, and do not use this medication for a continuing period.**

Consultation **1.**The following persons should consult a physician, pharmacist, or registered salesperson before taking this medication: (1) persons who are receiving any treatment from a physician, (2) women who are or may be pregnant, and (3) women who are breast-feeding.

2. If any of the symptoms mentioned below develop after taking ESTARON MOCHA ORAL SOLUTION, they might be side effects of the medication. Stop taking this medication immediately and consult a physician, pharmacist, or registered salesperson, presenting this bottle.

Gastrointestinal: inappetence and nausea/vomiting

Neuropsychiatric: tremor, dizziness, anxiety, sleeplessness, and headache

Cardiovascular: palpitations



[Indications] Elimination of sleepiness and malaise

[Dosage and Administration]

Adults (aged 15 years or older): Take a dose (30 mL per bottle) of ESTARON MOCHA ORAL SOLUTION once daily.

< Precautions Concerning Dosage and Administration >

(1) ESTARON MOCHA ORAL SOLUTION must not be taken by children under 15 years of age.

(2) This product must be taken strictly in accordance with the specified dosage and administration.

(3) This product is intended for once-daily administration. Do not take this product twice or more times in one day.

[Ingredients] Per bottle (30 mL)

Caffeine hydrate, 150 mg; thiamine chloride hydrochloride, 10 mg; pyridoxine hydrochloride, 5 mg; calcium glycerophosphate, 20 mg; nicotinamide, 15 mg; taurine, 1,000 mg

Excipients: D-sorbitol, white soft sugar, alcohol, propylene glycol, citric acid, sodium benzoate, paraben, ethyl vanillin, caramel, sodium L-glutamate, vanillin, and flavoring

[Precautions for Storage and Handling]

- (1) Store in a cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not take this medication after the expiration date.

Expiration date

Manufacturing No.



Relief System for Sufferers of
Adverse Drug Reactions
Phone: 0120-149-931

30mL

sleepiness and malaise

Medicine

Coffee-flavored solution to eliminate sleepiness

ESTARON MOCHA® ORAL SOLUTION

Class 3 OTC drug

Marketed by  SSP Co., Ltd.
3-20-2 Nishi-Shinjuku, Shinjuku-ku, Tokyo
Customer Service Office Phone: 0120-028-193
Opening hours: 9:00 to 17:00 (closed on Saturdays, Sundays, and Japanese public holidays)
<https://www.ssp.co.jp/>

Manufactured and Distributed by: Dojin Iyaku-Kako Co., Ltd.
8-1 Higashi-nakakawara, Senoue-machi, Fukushima-shi, Fukushima