

# MEMO<sup>®</sup> A

Class 2 OTC drug

## Ointment for traumatic injury

- **MEMO A** is a smooth and easy-to-spread ointment that contains a disinfectant, a topical anesthetic for relieving traumatic pain, and other ingredients to accelerate wound healing.
- The active ingredients of this product exert excellent effects on cuts, abrasions, and burns.
- MEMO A can be used as a home medication or a medication convenient for carrying when playing sports or traveling.



## Precautions for Use

### **Contraindications**

( If you do not follow the precautions exactly as listed below, you may be placed at increased risk of side effects. )

#### ① **The following persons must not use this medication:**

Persons who have had an allergic reaction to any of the ingredients of MEMO A or to chlorhexidine



### **Consultation**

#### ① **The following persons should consult a physician, pharmacist, or registered salesperson before using this medication:**

- (1) Persons who are receiving any treatment from a physician
- (2) Persons who have had any allergic reactions to drugs or other substances
- (3) Persons with lesions involving an extensive area of the skin
- (4) Persons with a deep wound or severe burn

#### ② **If any of the symptoms listed below develop after using MEMO A, they might be side effects of the medication. Stop using this medication immediately and consult a physician, pharmacist, or registered salesperson, presenting this leaflet.**

Affected area	Symptoms
Skin	Rash/redness and itching

In rare cases, any of the serious symptoms listed below may occur. In such cases, immediately consult a physician.

Name of the symptoms	Symptoms
Shock (anaphylaxis)	Itchy skin, urticaria, hoarse voice, sneezing, itchy throat, respiratory distress, palpitations, clouding of consciousness, etc., occurring immediately after using this medication

#### ③ **If the symptoms for which this medication is being used are not alleviated after applying the recommended doses for 5 or 6 days, stop using the medication and consult a physician, pharmacist, or registered salesperson, presenting this leaflet.**

## Indications

- Cuts, abrasions including shoe abrasions, puncture wounds, or bites and stings
- Slight burns
- Disinfection and antiseptics of a wound surface

## Dosage and Administration

Apply an appropriate amount of the ointment to the affected area several times daily.

### < Precautions Concerning Dosage and Administration >

- (1) Children must use this medication only under the supervision of their parents/guardians.
- (2) The ointment must be applied with caution so as not to come into contact with the eyes. Should it enter an eye, immediately wash the eye with cold or lukewarm water. If any severe reaction in the eye continues after washing, consult an ophthalmologist.
- (3) This medication is for external use only.

## Ingredients

Per gram

Chlorhexidine gluconate solution	10 mg
Dibucaine hydrochloride	3 mg
Allantoin	10 mg
Tocopherol acetate (vitamin E acetate)	1 mg
Zinc oxide	50 mg

Excipients: paraffin, petrolatum, white beeswax, sorbitan trioleate, phenol, thymol, and flavoring

## Treatment of cuts, abrasions, and slight burns...

Early treatment is important even for minor wounds such as cuts, abrasions, and slight burns. For the treatment of such injuries, MEMO A should be applied while paying attention to the following points:

- If sand or pebbles are attached to the wound surface, clean the wound area by washing with water, dry the area, and then apply **MEMO A**.
- In the case of a burn, immediately immerse the affected area in cold water for cooling and then apply **MEMO A**.
- Protect the affected area with an adhesive tape, gauze, etc., until the open wound has closed.
- Do not forcefully remove a scab formed on the wound surface.

## Precautions for Storage and Handling

- (1) Store the product in the can with the lid closed tightly, in a cool place away from direct sunlight. Closing the lid too tightly may lead to difficulty opening.
- (2) Keep out of the reach of children.
- (3) Do not transfer the contents to another container (which may lead to drug misuse or changes in the quality of this product).
- (4) Do not use this medication after the expiration date.

### Contact Information

**Inquiries should be made to the drugstore from where you purchased the product or to the customer service office.**

Customer Service Office, SSP Co., Ltd. Phone: 0120-028-193  
Opening hours: 9:00 to 17:00 (closed on Saturdays, Sundays, and Japanese public holidays)

Manufactured and  
Distributed by:



**SSP Co., Ltd.**

3-20-2 Nishi-Shinjuku, Shinjuku-ku, Tokyo 163-1444  
<https://www.ssp.co.jp/>

7-0AO

91