

NEW S.TAC® GRANULES

Class (2) OTC drug

Medication for common cold, an easy-to-swallow and easy-to-dissolve granule-type formulation that contains kakkonto-ka-kikyo extract

- **NEW S.TAC GRANULES** are a medication for common cold that are effective against symptoms such as chills, headache, sneezing, and nasal discharge. They contain Western medicine-derived drugs that are effective against various symptoms of common cold, as well as "kakkonto-ka-kikyo extract," a combination of kakkonto and kikyo, which has been used as a traditional herbal medicine for "common cold."
- **NEW S.TAC GRANULES** are easy-to-swallow and easy-to-dissolve granules.



Precautions for Use



Contraindications

(If you do not follow the precautions exactly as listed below, you may be placed at an increased risk of side effects or of having some accidents.)

● **The following persons must not take this medication:**

- (1) Persons who have had an allergic reaction to any of the ingredients of NEW S.TAC GRANULES
- (2) Persons who have experienced asthma after taking this medication, other common cold medications, or antipyretic-analgesics
- (3) Children under 12 years of age

● **Do not use any of the following medications during the period of taking NEW S.TAC GRANULES:**

Other common cold medications, antipyretic-analgesic medications, sedatives, antitussive-expectorant medications, or oral medications containing antihistamines (e.g., oral rhinitis medications, motion sickness medications, anti-allergic medications, or hypnotic sedatives)

● **Do not operate vehicles or machinery after taking this medication.**

(Sleepiness or drowsiness may occur.)

● **Breast-feeding women are advised to not take NEW S.TAC GRANULES or stop breast-feeding if this medication is to be taken.**

● **This medication must not be taken immediately before or after alcohol intake.**

● **Do not use this medication over a long continuing period.**



Consultation

● **The following persons should consult a physician, pharmacist, or registered salesperson before taking this medication:**

- (1) Persons who are receiving any treatment from a physician or dentist
- (2) Women who are or may be pregnant
- (3) Older persons
- (4) Persons who have had any allergic reactions to drugs or other substances
- (5) Persons with the following symptom: Hyperthermia (high fever) and dysuria
- (6) Persons who have received a diagnosis of any of the following diseases: thyroid dysfunction, diabetes mellitus, heart disease, hypertension, liver disease, kidney disease, gastroduodenal ulcers, glaucoma, respiratory dysfunction, obstructive sleep apnea syndrome, obesity, or phenylketonuria

● **If any of the symptoms listed below develop after taking NEW S.TAC GRANULES, they might be side effects of the medication. Stop taking this medication immediately and consult a physician, pharmacist, or registered salesperson, presenting this leaflet.**

Affected area	Symptoms
Skin	Rash/redness and itching
Gastrointestinal	Nausea/vomiting and inappetence
Neuropsychiatric	Dizziness

Affected area	Symptoms
Urinary	Dysuria
Other	Excessively decreased body temperature

In rare cases, any of the serious symptoms listed below may occur. In such cases, immediately consult a physician.

Name of the symptoms	Symptoms
Shock (anaphylaxis)	Itchy skin, urticaria, hoarse voice, sneezing, itchy throat, respiratory distress, palpitations, clouding of consciousness, etc., occurring immediately after taking this medication
Oculomucocutaneous syndrome (Stevens-Johnson syndrome), toxic epidermal necrolysis, or acute exanthematous pustulosis	Persistence or worsening of the following symptoms: high fever, ocular hyperemia, eye discharge, lip erosion, sore throat, rashes/redness over an extensive area of the skin, small rashes (small pustules) appearing on reddened skin, general malaise, or inappetence
Drug-induced hypersensitivity syndrome	Occurrence of symptoms such as redness over an extensive area of the skin, generalized exanthema, fever, malaise, or swelling of the lymph nodes (cervical, axillary, and/or inguinal lymph nodes)

(Continued from the front side)

Name of the symptoms	Symptoms
Hepatic impairment	Occurrence of fever, itching, rash, jaundice (yellowing of the skin and whites of the eyes), brown urine, general malaise, inappetence, etc.
Renal disorders	Occurrence of fever, rashes, decreased urine output, generalized edema/swelling, general malaise, arthralgia (pain in joints), diarrhea, etc.
Interstitial pneumonia	Sudden occurrence or persistence of the following symptoms: shortness of breath or respiratory distress when climbing stairs or slightly overworking, dry cough, or fever
Asthma	Occurrence of breathing sounds, such as wheezing and whistling, or respiratory distress
Aplastic anemia	The following symptoms can occur: bruising, nasal bleeding, gingival bleeding, fever, pallor of the skin or mucous membranes, fatigue, palpitations, shortness of breath, feeling sick and fainting, and hematuria.
Agranulocytosis	Sudden high fever, chills, sore throat, etc.
Respiratory depression	Occurrence of shortness of breath, respiratory distress, etc.

- Any of the conditions mentioned below may occur after taking NEW S.TAC GRANULES. If such symptoms persist or worsen, stop taking this medication and consult a physician, pharmacist, or registered salesperson, presenting this leaflet.
Constipation, thirst, or sleepiness
- If the symptoms of common cold do not alleviate after taking 5 or 6 doses, stop taking this medication and consult a physician, pharmacist, or registered salesperson, presenting this leaflet.

Indications Alleviation of various symptoms of common cold (fever, chills, headache, nasal discharge, nasal stuffiness, sneezing, cough, sore throat, joint pain, and muscle pain)

Dosage and Administration Take a dose of NEW S.TAC GRANULES, according to the table below, with cold or lukewarm water, 3 times daily. Each dose should be taken within 30 minutes after a meal, whenever possible.

Age	Adults (aged 15 years or older)	Children aged 12 to 14 years	Children under 12 years of age
Dose	1 sachet	2/3 sachets (two-thirds of one sachet)	✗ This medication must not be taken.

< Precautions Concerning Dosage and Administration >

- (1) NEW S.TAC GRANULES must be used strictly in accordance with the specified dosage and administration.
- (2) Children must take this product only under the supervision of their parents/guardians.

Ingredients

per 1 sachet
(2.0 g)

Kakkonto-ka-kikyo extract... 1,000 mg

Extract prepared from the decoction of 0.53 g of pueraria root, 0.27 g of ephedra herb, 0.2 g of cinnamon bark, 0.2 g of peony root, 0.27 g of jujube, 0.07 g of ginger, 0.13 g of glycyrrhiza, and 0.27 g of platycodon root

Acetaminophen 240 mg
Dihydrocodeine phosphate 5 mg
Chlorpheniramine maleate 2.5 mg
Anhydrous caffeine 25 mg

Excipients:carmellose calcium, silicic anhydride, gelatin, lactose, white soft sugar, starch, corn starch, aspartame (L-phenylalanine compound), vanillin, and flavoring

Precautions for Storage and Handling

- (1) Store in a dry, cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer the contents to another container (which may lead to drug misuse or changes in the quality of this product).
- (4) When the granules in one sachet are to be taken in divided doses, the remaining granules must be kept in the sachet with its opening portion folded down, and the contents must be used within 2 days.
- (5) Do not take this medication after the expiration date.

Contact Information	Inquiries should be made to the drugstore from where you purchased the product or to the customer service office. Customer Service Office, SSP Co., Ltd. Phone: 0120-028-193 Opening hours: 9:00 to 17:00 (closed on Saturdays, Sundays, and Japanese public holidays)
----------------------------	---



Manufactured and Distributed by:
SSP Co., Ltd.

3-20-2 Nishi-Shinjuku, Shinjuku-ku, Tokyo 163-1444
<https://www.ssp.co.jp/>