

この説明書は製品についての参考情報です。正しくは製品に添付されている説明書をご確認ください。

This document serves as reference information. Please ensure to read the document attached to the product.

Underlined parts: revised in July 2019

[Keep this information leaflet with this medication and be sure to read it before taking the medication.]

# SS BRON® TABLETS

Class ② OTC drug

Antitussives-expectorant medication < sugar-coated tablets > for cough and sputum

- Cough is a symptom associated with unhealthy conditions of the throat and bronchus. Persistent cough burdens the lungs and/or heart and interferes with sleep, leading to physical wasting.
- **SS BRON TABLETS** are white sugar-coated tablets that are effective in suppressing cough and promoting sputum expectoration.



## Precautions for Use



### Contraindications

(If you do not follow the precautions exactly as listed below, your current symptoms may worsen, and you may be placed at increased risk of side effects or accidents.)

#### ① The following persons must not take this medication:

Children under 12 years of age

#### ② Do not use any of the following medications during the period of taking SS BRON TABLETS:

Other antitussive-expectorant medications, common cold medications, sedatives, or oral medications containing antihistamines (e.g., oral rhinitis medications, motion sickness medications, anti-allergic medications, or hypnotic sedatives)

#### ③ Do not operate vehicles or machinery after taking this medication.

(Sleepiness or drowsiness may occur.)

#### ④ Breast-feeding women are advised to not take SS BRON TABLETS or stop breast-feeding if this medication is to be taken.

#### ⑤ Do not take more than the specified dosage of this medication or use it over a long continuing period.



### Consultation

#### ① The following persons should consult a physician, pharmacist, or registered salesperson before taking this medication:

- (1) Persons who are receiving any treatment from a physician
- (2) Women who are or may be pregnant
- (3) Older persons
- (4) Persons who have had any allergic reactions to drugs or other substances
- (5) Persons with the following symptom: Hyperthermia (high fever) and dysuria
- (6) Persons who have received a diagnosis of any of the following diseases: heart disease, hypertension, diabetes mellitus, glaucoma, thyroid dysfunction, respiratory dysfunction, obstructive sleep apnea syndrome, or obesity

#### ② If any of the symptoms listed below develop after taking SS BRON TABLETS, they might be side effects of the medication. Stop taking this medication immediately and consult a physician, pharmacist, or registered salesperson, presenting this leaflet.

Affected area	Symptoms
Skin	Rash/redness and itching
Gastrointestinal	Nausea/vomiting and inappetence
Neuropsychiatric	Dizziness
Urinary	Dysuria

In rare cases, any of the serious symptoms listed below may occur. In such cases, immediately consult a physician.

Name of the symptoms	Symptoms
Aplastic anemia	The following symptoms can occur: bruising, nasal bleeding, gingival bleeding, fever, pallor of the skin or mucous membranes, fatigue, palpitations, shortness of breath, feeling sick and fainting, and hematuria.
Agranulocytosis	Sudden high fever, chills, sore throat, etc.
Respiratory depression	Occurrence of shortness of breath, respiratory distress, etc.

#### ③ Any of the conditions mentioned below may occur after taking SS BRON TABLETS. If such symptoms persist or worsen, stop taking this medication and consult a physician, pharmacist, or registered salesperson, presenting this leaflet. Constipation, thirst, or sleepiness

#### ④ If the symptoms for which this medication is being taken are not alleviated after 5 or 6 doses, stop taking this medication and consult a physician, pharmacist, or registered salesperson, presenting this leaflet.

**Indications**

Cough and sputum

**Dosage and Administration**

Take a dose of SS BRON TABLETS with cold or lukewarm water thrice daily at intervals of not less than 4 hours, according to the table below.

Age	Adults (aged 15 years or older)	Children aged 12 to 14 years	Children under 12 years of age
Dose	4 tablets	2 tablets	× This medication must not be taken.

**< Precautions Concerning Dosage and Administration >**

- (1) SS BRON TABLETS must be used strictly in accordance with the specified dosage and administration.
- (2) Children must take this product only under the supervision of their parents/guardians.

**Ingredients and Actions**

In a total of 12 tablets (daily dose for adults)

Ingredients	Amount per 12 tablets	Action
Dihydrocodeine phosphate	30 mg	To inhibit the cough reflex by acting on the cough center in the medulla oblongata.
dl-Methylephedrine hydrochloride	50 mg	To relieve bronchial muscle tone and alleviate cough, promoting sputum expectoration
Chlorpheniramine maleate	8 mg	To alleviate allergy cough
Anhydrous caffeine	90 mg	To assist the activities of the other ingredients

Excipients: carmellose calcium, silicic anhydride, cellulose, lactose, white soft sugar, hydroxypropylcellulose, hypromellose, povidone, calcium carbonate, carnauba wax, magnesium stearate, talc, titanium oxide, pullulan, and polyoxyethylene polyoxypropylene glycol


**For the treatment of cough and sputum...**

**In addition to taking this medication, attention should be paid to the following points:**

- Try to restore physical strength by having sufficient sleep and following a nutritious diet.
- Make sure indoor air does not become overly dry.
- Pay attention to throat care such as by gargling.

**Precautions for Storage and Handling**

- (1) Store the product in the jar with the lid closed tightly, in a dry, cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer the contents to another container (which may lead to drug misuse or changes in the quality of this product).
- (4) Do not handle this product with wet hands. If the product is handled with wet hands, the sugar coating may become uneven or discolored.
- (5) The filler in the jar is intended to prevent breakage of tablets during transportation. After the package has been opened, the filler is no longer necessary and must be removed.
- (6) Do not take this medication after the expiration date.

**Contact Information**

**Inquiries should be made to the drugstore from where you purchased the product or to the customer service office.**

Customer Service Office, SSP Co., Ltd. Phone: 0120-028-193  
Opening hours: 9:00 to 17:00 (closed on Saturdays, Sundays, and Japanese public holidays)



Manufactured and Distributed by:

**SSP Co., Ltd.**

3-20-2 Nishi-Shinjuku, Shinjuku-ku, Tokyo 163-1444

<https://www.ssp.co.jp/>