

この説明書は製品についての参考情報です。正しくは製品に添付されている説明書をご確認ください。
This document serves as reference information. Please ensure to read the document attached to the product.

[Keep this information leaflet with this medication and be sure to read it before taking the medication.]

SS BRON[®] SOLUTION L

Class 2 OTC
drug

Antitussives-expectorant medication for cough and sputum

- Persistent cough not only is troublesome but also prevents sound sleep and can lead to physical wasting.
Sputum stuck in the throat is also very distressing.
- **SS BRON SOLUTION L** is an easy-to-swallow syrup that is very effective against cough and sputum. It contains ingredients to suppress cough and promote sputum expectoration.

Precautions for Use



Contraindications

(If you do not follow the precautions exactly as listed below, your current symptoms may worsen, and you may be placed at increased risk of side effects or accidents.)

① The following persons must not take this medication:

Persons who have had an allergic reaction to any of the ingredients of SS BRON SOLUTION L

② Do not take any of the following medications during the period of taking SS BRON SOLUTION L:

Other antitussive-expectorant medications, common cold medications, sedatives, or oral medications containing antihistamines (e.g., oral rhinitis medications, motion sickness medications, anti-allergic medications, or hypnotic sedatives)

③ Do not operate vehicles or machinery after taking this medication.

(Sleepiness or drowsiness may occur.)



Consultation

① The following persons should consult a physician, pharmacist, or registered salesperson before taking this medication:

- (1) Persons who are receiving any treatment from a physician
- (2) Women who are or may be pregnant
- (3) Persons who have had any allergic reactions to drugs or other substances
- (4) Persons with the following symptom: Hyperthermia (high fever) and dysuria
- (5) Persons who have received a diagnosis of the following disease: glaucoma



② If any of the symptoms listed below develop after taking SS BRON SOLUTION L, they might be side effects of the medication. Stop taking this medication immediately and consult a physician, pharmacist, or registered salesperson, presenting this leaflet.

Affected area	Symptoms
Skin	Rash/redness and itching
Gastrointestinal	Nausea/vomiting and inappetence
Neuropsychiatric	Dizziness
Respiratory	Respiratory distress or shortness of breath
Urinary	Dysuria

In rare cases, any of the serious symptoms listed below may occur. In such cases, immediately consult a physician.

Name of the symptoms	Symptoms
Shock (anaphylaxis)	Itchy skin, urticaria, hoarse voice, sneezing, itchy throat, respiratory distress, palpitations, clouding of consciousness, etc., occurring immediately after taking this medication
Aplastic anemia	The following symptoms can occur: bruising, nasal bleeding, gingival bleeding, fever, pallor of the skin or mucous membranes, fatigue, palpitations, shortness of breath, feeling sick and fainting, and hematuria.
Agranulocytosis	Sudden high fever, chills, sore throat, etc.

③ Any of the conditions mentioned below may occur after taking SS BRON SOLUTION L. If such symptoms persist or worsen, stop taking this medication and consult a physician, pharmacist, or registered salesperson, presenting this leaflet.

Thirst and sleepiness

④ If the symptoms for which this medication is being taken are not alleviated after 5 or 6 doses, stop taking this medication and consult a physician, pharmacist, or registered salesperson, presenting this leaflet.

Indications

Cough and sputum

Dosage and Administration

Take a dose of SS BRON SOLUTION L using the accompanying measuring cup thrice daily after meals, according to the table below. This medication can be taken up to 6 times a day as needed, at intervals of not less than 4 hours.

Age	Adults (aged 15 years or older)	Children aged 11 to 14 years	Children aged 8 to 10 years	Children aged under 8 years
Dose	5 mL	3.3 mL	2.5 mL	× This medication must not be taken.

* Keep the measuring cup clean by washing with water after each use, and store it with this medication.

< Precautions Concerning Dosage and Administration >

- (1) SS BRON SOLUTION L must be used strictly in accordance with the specified dosage and administration.
- (2) Children must take this product only under the supervision of their parents/guardians.

Ingredients and Actions

Ingredient	Amount per 30 mL	Action
Dextromethorphan hydrobromide hydrate	60 mg	To inhibit the cough reflex by acting on the cough center located in the medulla oblongata, which controls coughing
Guaifenesin	170 mg	To enhance the function of respiratory mucosal secretion and reduce the viscosity of sputum, promoting sputum expectoration
Chlorpheniramine maleate	12 mg	To alleviate allergy cough
Anhydrous caffeine	62 mg	To assist the activities of the other ingredients

Excipients: high-fructose corn syrup (HFCS), alcohol, propylene glycol, sodium chloride, citric acid, sodium benzoate, paraben, ethyl vanillin, caramel, vanillin, l-menthol, and flavoring


For the treatment of cough and sputum...

In addition to taking this medication, attention should be paid to the following points:

- Try to restore physical strength by having sufficient sleep and following a nutritious diet.
- Make sure indoor air does not become overly dry.
- Pay attention to throat care such as by gargling.

Precautions for Storage and Handling

- (1) Store the product in the bottle with the cap closed tightly, in a cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer the contents to another container (which may lead to drug misuse or changes in the quality of this product).
- (4) Before replacing the cap, wipe the mouth of the bottle thoroughly. If the cap is closed with some drug solution remaining on the mouth of the bottle, this may cause the cap to not open easily at the next use. If the cap is not tightly closed, this may cause leakage or contamination of the solution.
- (5) Do not take this medication after the expiration date.



Contact Information

Inquiries should be made to the drugstore from where you purchased the product or to the customer service office.

Customer Service Office, SSP Co., Ltd. Phone: 0120-028-193

Opening hours: 9:00 to 17:00 (closed on Saturdays, Sundays, and Japanese public holidays)



Manufactured and Distributed by:

SSP Co., Ltd.

3-20-2 Nishi-Shinjuku, Shinjuku-ku, Tokyo 163-1444

<https://www.ssp.co.jp/>