

00007772

この説明書は製品についての参考情報です。正しくは製品に添付されている説明書をご確認ください。
This document serves as reference information. Please ensure to read the document attached to the product.

[Keep this information leaflet with this medication and be sure to read it before taking the medication.]

BUSCOPAN[®]-A TABLETS

Class 2 OTC drug

Analgesic antispasmodic agent for stomachache and abdominal pain

- Pain, including stomachache, abdominal pain, and stomach cramps, occurs due to excessively increased gastrointestinal tone (hypertonia) or gastrointestinal cramps.
- **BUSCOPAN-A TABLETS** are effective for relieving abnormal gastrointestinal tone (hypertonia) and pain including stomachache, abdominal pain, and stomach cramps.

⚠ Precautions for Use



Contraindications

If you do not follow the precautions exactly as listed below, your current symptoms may worsen, and you may be placed at increased risk of side effects or accidents.

- ① **The following persons must not take this medication:**
Persons who have had an allergic reaction to any of the ingredients of BUSCOPAN-A TABLETS.
- ② **Do not take any of the following medications during the period of taking BUSCOPAN-A TABLETS:**
Other gastrointestinal antispasmodic medications, other gastrointestinal medications containing scopolia extract, and motion sickness medications
- ③ **Do not operate vehicles or machinery after taking this medication.**
(Symptoms such as blurred vision and abnormal brightness may occur.)



Consultation

- ① **The following persons should consult a physician, pharmacist, or registered salesperson before taking this medication:**
 - (1) Persons who are receiving any treatment from a physician
 - (2) Women who are or may be pregnant
 - (3) Older persons
 - (4) Persons who have had any allergic reactions to drugs or other substances
 - (5) Persons with the following symptom: Dysuria
 - (6) Persons who have received a diagnosis of any of the following diseases: Heart disease and glaucoma
- ② **If any of the symptoms listed below develop after taking BUSCOPAN-A TABLETS, they might be side effects of the medication. Stop taking this medication immediately and consult a physician, pharmacist, or registered salesperson, presenting this leaflet.**



Affected area	Symptoms
Skin	Rash/redness and itching
Neuropsychiatric	Headache
Urinary	Dysuria
Other	Facial hot flushes and abnormal sensitivity to light

In rare cases, any of the serious symptoms listed below may occur. In such cases, immediately consult a physician.

Name of the symptoms	Symptoms
Shock (Anaphylaxis)	Itchy skin, urticaria, hoarse voice, sneezing, itchy throat, respiratory distress, palpitations, clouding of consciousness, etc., occurring immediately after taking this medication

- ③ **Any of the conditions mentioned below may occur after taking BUSCOPAN-A TABLETS. If such symptoms persist or worsen, stop taking this medication and consult a physician, pharmacist, or registered salesperson, presenting this leaflet.**
Thirst, constipation, and blurred vision
- ④ **If gastrointestinal symptoms are not alleviated after taking 5 or 6 doses, stop taking this medication and consult a physician, pharmacist, or registered salesperson, presenting this leaflet.**

Indications

Stomachache, abdominal pain, stomach cramps (abdominal colic and spasms), hyperchlorhydria (excessive gastric acid secretion), and heartburn



Stomachache



Abdominal pain

Dosage and Administration

Take a dose of BUSCOPAN-A TABLETS with cold or lukewarm water up to thrice daily at intervals of not less than 4 hours, according to the table below.

Age	Adults (aged 15 years or older)	Children under 15 years of age
Dose	1 tablet	× This medication must not be taken.

< Precautions Concerning Dosage and Administration >

- (1) BUSCOPAN-A TABLETS must be used strictly in accordance with the specified dosage and administration.
- (2) How to remove the tablet: As shown in the illustration on the right, firmly press the raised portion of the press-through package (PTP) with a fingertip to push the tablet through the aluminum foil on the reverse side. (If the PTP sheet is mistakenly swallowed, its sharp edges may perforate the esophageal mucosa or lead to incidental events.)



Ingredients


Scopolamine butylbromide..... 10 mg

Per tablet
(dose per administration)

Excipients: lactose, white soft sugar, macrogol, acacia, carnauba wax, white beeswax, magnesium stearate, shellac, talc, titanium oxide, corn starch, sodium lauryl sulfate, and tartaric acid

Precautions for Storage and Handling

- (1) Store in a dry, cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer the contents to another container (which may lead to drug misuse or changes in the quality of this product).
- (4) Do not take this medication after the expiration date.



Contact Information

Inquiries should be made to the drugstore from where you purchased the product or to the customer service office.
 Customer Service Office, SSP Co., Ltd. Phone: 0120-028-193
 Opening hours: 9:00 to 17:00 (closed on Saturdays, Sundays, and Japanese public holidays)



Manufactured and Distributed by:

SSP Co., Ltd.

3-20-2 Nishi-Shinjuku, Shinjuku-ku, Tokyo 163-1444
<https://www.ssp.co.jp/>