


この説明書は製品についての参考情報です。正しくは製品のパッケージをご確認ください。パッケージデザインは変更されている可能性があります。

This document serves as reference information. Please ensure to read the information on the package.

The package design may have been updated.



Be sure to read the instructions given on the inside of the carton before use. Open the carton by separating the overlapped portion along the perforations.

NEW BRON SOLUTION - ACE

©: Registered trademark

(1) Press here to open.

Then, press the portion marked with (2) to open further.

For cough and sputum

Class ② OTC drug

NEW BRON® SOLUTION - ACE

120mL

Paper Carton Plastic Cap: PP Cup wrapping: PP Outer packaging film: PE and PP

Manufactured and Distributed by:
SSP Co., Ltd.
3-20-2 Nishi-Shinjuku, Shinjuku-ku, Tokyo
Customer Service Office Phone: 0120-028-193
Opening hours: 9:00 to 17:00 (closed on Saturdays, Sundays, and Japanese public holidays)
<https://www.ssp.co.jp/>

Expiration date

- [Caution] 1. Do not take more than the specified dosage of this medication or use it over a long continuous period.
2. NEW BRON SOLUTION - ACE must be taken correctly according to the specified dosage and administration.
3. The following persons must not take this medication: Children under 12 years of age.
4. Do not operate vehicles or machinery after taking this medication.
5. Breast-feeding women are advised to not take NEW BRON SOLUTION - ACE or stop breast-feeding if this medication is to be taken.
6. The following persons should consult a physician, pharmacist, or registered salesperson before taking this medication:
(1) Persons who are receiving any treatment from a physician
(2) Women who are or may be pregnant
(3) Persons who have had any allergic reactions to drugs or other substances
(4) Persons with the following symptom: Hyperthermia (high fever) and dysuria
(5) Persons who have received a diagnosis of any of the following diseases: Glaucoma, respiratory dysfunction, obstructive sleep apnea syndrome, or obesity
7. Carefully read the instructions given on the inside of the carton before taking this product.
8. Store the product in the bottle with the cap closed tightly, in a cool place away from direct sunlight.
9. After each use, wipe the mouth of the bottle thoroughly before replacing the cap.

For cough and sputum

[Indications] Cough and sputum
[Dosage and Administration] Take a dose of NEW BRON SOLUTION - ACE using the accompanying measuring cup thrice daily after meals, according to the table below. This medication can be taken up to 6 times a day as needed; however, if taken more than 3 times a day, the medication must be taken at intervals of not less than 4 hours.

Age	Adults (aged 15 years or older)	Children aged 12 to 14 years
Dose	10mL	6.6mL

[Ingredients] per 60 mL
Dihydrocodeine phosphate 30 mg
Guaifenesin 170 mg
Chlorpheniramine maleate 12 mg
Anhydrous caffeine 62 mg

Excipients: high-fructose corn syrup (HFCS), alcohol, propylene glycol, sodium chloride, citric acid, sodium benzoate, paraben, ethyl vanillin, caramel, vanillin, l-menthol, and flavoring

NEW BRON SOLUTION - ACE

NEW BRON SOLUTION - ACE

NEW BRON SOLUTION - ACE

Relief System for Sufferers of Adverse Drug Reactions Phone: 0120-149-931

NEW BRON SOLUTION-ACE

ANTITUSSIVE & EXPECTORANT

NEW BRON SOLUTION-ACE

ANTITUSSIVE & EXPECTORANT

O

Then, separate the portion marked with (3) from the upper side.

(2) Press here to open.

(4)

SSP CO., LTD. TOKYO JAPAN

Class ② OTC drug



Manufactured and Distributed by:
SSP Co., Ltd.

Class ② OTC drug

SSP CO., LTD. TOKYO JAPAN

Class ② OTC drug



To protect the environment, no information leaflet is co-packaged.

Keep this carton in a safe place and be sure to read the instructions given on the carton before use.

Class ② OTC drug

NEW BRON[®] SOLUTION - ACE

⚠ Precautions for Use

⊗ Contraindications

(If you do not follow the precautions exactly as listed below, your current symptoms may worsen, and you may be placed at increased risk of side effects or accidents.)

- ❶ **The following persons must not take this medication:**
Children under 12 years of age
- ❷ **Do not use any of the following medications during the period of taking NEW BRON SOLUTION - ACE:**
Other antitussive-expectorant medications, common cold medications, sedatives, and oral medications containing antihistamines (e.g., oral rhinitis medications, motion sickness medications, anti-allergic medications, and hypnotic sedatives)
- ❸ **Do not operate vehicles or machinery after taking this medication.**
(Sleepiness or drowsiness may occur.)
- ❹ **Breast-feeding women are advised to not take NEW BRON SOLUTION - ACE or stop breast-feeding if this medication is to be taken.**
- ❺ **Do not take more than the specified dosage of this medication or use it over a long continuous period.**
(Malaise, collapse, etc., may occur.)

👤 Consultation

- ❶ **The following persons should consult a physician, pharmacist, or registered salesperson before taking this medication:**
 - (1) Persons who are receiving any treatment from a physician
 - (2) Women who are or may be pregnant
 - (3) Persons who have had any allergic reactions to drugs or other substances
 - (4) Persons with the following symptom: Hyperthermia (high fever) and dysuria
 - (5) Persons who have received a diagnosis of any of the following diseases: Glaucoma, respiratory dysfunction, obstructive sleep apnea syndrome, or obesity
- ❷ **If any of the symptoms listed below develop after taking NEW BRON SOLUTION - ACE, they might be side effects of the medication. Stop taking this medication immediately and consult a physician, pharmacist, or registered salesperson, presenting this carton.**

Affected area	Symptoms
Skin	Rash/redness and itching
Gastrointestinal	Nausea/vomiting and inappetence
Neuropsychiatric	Dizziness
Urinary	Dysuria

In rare cases, any of the serious symptoms listed below may occur. In such cases, immediately consult a physician.

Name of the symptoms	Symptoms
Aplastic anemia	The following symptoms can occur: bruising, nasal bleeding, gingival bleeding, fever, pallor of the skin or mucous membranes, fatigue, palpitations, shortness of breath, feeling sick and fainting, and hematuria.
Agranulocytosis	Occurrence of sudden high fever, chills, sore throat, etc.
Respiratory depression	Occurrence of shortness of breath, respiratory distress, etc.

- ❸ **Any of the conditions mentioned below may occur after taking NEW BRON SOLUTION - ACE. If such symptoms persist or worsen, stop taking this medication and consult a physician, pharmacist, or registered salesperson, presenting this carton.** Constipation, thirst, and sleepiness
- ❹ **If the symptoms for which this medication is being taken are not alleviated after 5 or 6 doses, stop taking this medication and consult a physician, pharmacist, or registered salesperson, presenting this carton.**

Indications

Cough and sputum

Dosage and Administration

Take a dose of NEW BRON SOLUTION - ACE using the accompanying measuring cup thrice daily after meals, according to the table below. This medication can be taken up to 6 times a day as needed; however, if taken more than 3 times a day, the medication must be taken at intervals of not less than 4 hours.

Age	Adults (aged 15 years or older)	Children aged 12 to 14 years	Children under 12 years of age
Dose	10 mL	6.6 mL	✗ This medication must not be taken.

* Keep the measuring cup clean by washing it with water after each use, and store it with this medication.

< Precautions Concerning Dosage and Administration >

- (1) NEW BRON SOLUTION - ACE must be used strictly in accordance with the specified dosage and administration.
- (2) Children must take this product only under the supervision of their parents/guardians.

Ingredients and Actions

Ingredients	Amount per 60 mL	Action
Dihydrocodeine phosphate	30 mg	To inhibit the cough reflex by acting on the cough center in the medulla oblongata
Guaifenesin	170 mg	To enhance the function of respiratory mucosal secretion and reduce the viscosity of sputum, promoting sputum expectoration
Chlorpheniramine maleate	12 mg	To alleviate allergy cough
Anhydrous caffeine	62 mg	To assist the activities of the other ingredients

Excipients: high-fructose corn syrup (HFCS), alcohol, propylene glycol, sodium chloride, citric acid, sodium benzoate, paraben, ethyl vanillin, caramel, vanillin, l-menthol, and flavoring

Precautions for Storage and Handling

- (1) Store the product in the bottle with the cap closed tightly, in a cool place away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer the contents to another container (which may lead to drug misuse or changes in the quality of this product).
- (4) Before replacing the cap, wipe the mouth of the bottle thoroughly. If the cap is closed with some drug solution attached to the mouth of the bottle, this may cause the cap to not open easily at the next use. If the cap is not tightly closed, this may cause leakage or contamination of the solution.
- (5) Do not take this medication after the expiration date.

Contact Information

Inquiries should be made to the drugstore from where you purchased the product or to the customer service office.
Customer Service Office, SSP Co., Ltd. Phone: 0120-028-193
Opening hours: 9:00 to 17:00 (closed on Saturdays, Sundays, and Japanese public holidays)

©: Registered trademark